A NEW SPECIES OF THE GENUS *EUCHORTHIPPUS* TARBINSKY FROM SHANDONG PROVINCE (ORTHOPTERA, ACRIDOIDEA, ARCYPTERIDAE)

YIN Xiang-Chu^{1,2}, XIAO Yun-Li¹

- 1. College of Life Sciences, Hebei University, Baoding 071002, China; E-mail; yxch@ sdau. edu. cn
- 2. College of Plant Protection, Shandong Agricultural University, Taian 271018, China

Abstract A new species of the genus *Euchorthippus* Tarbinsky, 1926, namely *E. shandongensis* sp. nov. from Shandong Province of China is described in this paper.

Euchorthippu shandongensis sp. nov. (Figs 1 - 8)

The new species is similar to Euchorthippus unicolor (Ikonnikov, 1913), but differs from the latter in: fastigial foveolae longer, length 3.2 times width; frontal ridge below the median ocellus narrow, downward broaden distinctly; tegmina of male longer, extending over base of epiproct distinctly; subgenital plate of male shorter, length 1.5 times width in lateral view; arc of epiphallus deep, lower apex with acute

tuber on both side. The type specimens are deposited in College of Plant Protection, Shandong Agricultural University, Taian, China.

Holotype δ , Kunyu Mountain, Shangdong (37°14′N, 121°46′E; alt. 230 – 923 m), 26 Sep. 2008. Paratypes: 4 δ δ , 2 \circ \circ , collected by XIAO Yun-Li, LI Xin-Jiang and YE Bao-Hua.

Length of body: $3 \cdot 15.0 - 16.5 \text{ mm}$, $9 \cdot 22.0 - 22.3 \text{ mm}$. Length of tegmen: $3 \cdot 9.0 - 9.8 \text{ mm}$, $9 \cdot 12.0 - 12.5 \text{ mm}$. Length of hind femur: $3 \cdot 10.9 - 11.2 \text{ mm}$, $9 \cdot 13.8 - 14.8 \text{ mm}$.

Etymology. The specific epithet is named for Shandong, the type locality,

Key words Orthoptera, Acridoidea, Arcypteridae, Euchorthippus, new species, China.

山东省异爪蝗属一新种 (直翅目,蝗总科,网翅蝗科)

印象初1,2 肖云丽1

- 1. 河北大学生命科学学院 保定 071002, E-mail: yxch@ sdau. edu. cn
- 2. 山东农业大学植物保护学院 泰安 271018

摘 要 记述采自中国山东省网翅蝗科异爪蝗属 Euchorthippus Tarbinsky, 1926 1 新种, 山东异爪蝗 Euchorthippus shandongensis sp. nov.。新种同素色异爪蝗 Euchorthippus unicolor (Ikonnikov, 1913) 近似, 其区别特征为: 头侧窝较长, 长为宽的3.2 倍; 颜面隆起明显, 具纵沟, 中眼之下缩狭, 往下明显宽大; 雄性前翅较长, 明显超过肛上板基部; 雄性下生殖板较短, 侧面观长为宽的1.5 倍; 阳茎基背片弓形深, 两侧下端具尖突。模式标本保存于山东农业大学植保学院。

关键词 直翅目,蝗总科,网翅蝗科,异爪蝗属,新种,中国. 中图分类号 Q969.265.1

异爪蝗属 Euchorthippus 于 1926 年由 Tarbinsky 建立,属网翅蝗科、网翅蝗亚科,迄今为止,全世界已知 30 种,其中 20 种分布在中国 [Fischer-Waldheim, 1846; Ikonnikov, 1913; Tarbinsky, 1925; Chang, 1937; Tu et al., 1964; Cheng et al., 1965; Zheng, 1980a; Zheng, 1980b; Jin et al., 1983; Zhang et al., 1985; Zheng, 1986; Bi et al., 1987; Zhang et al., 1992a; Zhang et al., 1992b; Liang et al.,

1992; Zheng, 1993; Zheng et al., 1993; Yin et al., 1996; Zheng et al., 1998; Ren et al., 2001; Zheng et al., 2002]。记述了于 2008 年采自山东省异爪蝗属 Euchorthippus Tarbinsky, 1926 1 新种, 山东异爪蝗 Euchorthippus shandongensis sp. nov., 记述如下。模式标本保存于山东农业大学植保学院标本室。

山东异爪蝗,新种 Euchorthippus shandongensis sp.

nov. (图 1~8)

雄性(图1~6) 体小型。头顶宽短,顶钝,复眼前缘头顶最大宽度为头顶长的1.5倍。颜面倾斜,颜面隆起明显,具纵沟,中眼之下缩狭,往下明显宽大。头侧窝长方形,长为宽的3.2倍。复眼椭圆型,其垂直直径为水平直径的2倍,为眼下沟长的2.4倍。触角丝状,细长,明显超过前胸背板的后缘,中段一节长为宽的2.3倍。前胸背板前缘较直,后缘呈弧形突出,中隆线明显,被后横沟割断,后横沟位于中部之后,沟前区的长度为沟后区长的1.2倍。前胸背板侧隆线在沟前区呈钝角形弯曲,在沟

后区向外开张。中胸腹板中隔长为最小宽度的 1.2 倍。后胸腹板侧叶分开较狭。前翅较长,明显超过肛上板基部,缘前脉域近基部明显扩大,顶端不超过前翅的中部;肘脉域的最大宽度大于中脉域的最大宽度。后足股节细长,长为最宽处的 5.6 倍。后足胫节外缘和内缘各具刺 12 个,缺外端刺。跗节爪间中垫大,到达爪的端部,两爪长度不等,外爪明显长于内爪。鼓膜孔半圆形。下生殖板短锥状,较短,侧面观长为宽的 1.5 倍,顶端钝圆。阳茎基背片猫状突端部略细,弓形深,两侧下端具尖突。



图 1~8 山东异爪蝗,新种 Euchorthippus shandongensis sp. nov. 1,3~4. ♂7.♀ 1. 侧面 (lateral view) 2. 头侧窝 (fastigial foveolae) 3. 颜面隆起 (frontal ridge) 4. 下生殖板 (subgenital plate) 5. 阳茎基背片 (epiphallus) 6. 爪 (claw) 7. 侧面 (lateral view) 8. 产卵瓣 (ovipositor valves)

雌性(图7~8) 近似雄性,体型较大。触角丝状,基部的触角节略宽大,较短,略超过前胸背板的后缘,中段一节长为宽的1.8倍。前翅较短,超过后足股节的中部。中胸腹板中隔长为最小宽度的1.1倍。下生殖板后缘中央呈弧形突出。产卵瓣细长,边缘光滑,顶端弯曲,呈钩状。

体通常暗褐色。前翅黄褐色。后足股节外侧黄褐色,内侧红褐色。后足胫节淡黄色。后足跗节淡黄色。

体长: \$15.0~16.5 mm, \$22.0~22.3 mm。 前翅长: \$9.0~9.8 mm, \$12.0~12.5 mm。后足 股节长: \$10.9~11.2 mm, \$13.8~14.8 mm。 正模 δ , 山东昆嵛山(37°14′N,121°46′E;海拔230~923 m),2008-09-26,肖云丽、李新江、叶保华采。副模: $4\delta\delta$, 2♀♀,记录同正模。模式标本保存于山东农业大学植保学院标本室。

新种同素色异爪蝗 Euchorthippus unicolor (Ikonnikov, 1913) 近似, 其区别特征见表 1。 词源: 新种种名源自新种模式标本产地。

表 1 山东异爪蝗和素色异爪蝗特征比较

Table 1. Comparison of Euchorthippus shandongensis sp. nov. and Euchorthippus unicolor.

	山东异爪蝗, 新种 Euchorthippus shandongensis sp. nov.	素色异爪蝗 Euchorthippus unicolor
头侧窝	长为宽的 3. 2 倍	长为宽的 2. 5 倍
Fastigial foveolae	Length is 3. 2 times width	Length is 2. 5 times width
颜面隆起	中眼之下缩狭,往下明显宽大	近乎平行
Frontal ridge	Below the median ocellus narrow, downward broaden distinctly	Nearly parallel
雄性前翅	较长,明显超过肛上板基部	较短,到达肛上板基部
Tegmina of male	Longer, extending over the base of epiproct distinctly	Shorter, reaching the base of epiproct
雄性下生殖板	较短, 侧面观长为宽的 1.5 倍	较长,侧面观长为宽的2.3倍
Subgenital plate of male	Shorter, length is 1.5 times width in lateral view	Longer, length 2.3 times width in lateral view
阳茎基背片	弓形深,两侧下端具尖突	弓形浅,两侧下端不具尖突
Epiphallus	Arc deep, lower apex with acute tuber on both side	Arc shallow, lower apex without acute tuber on both side

致谢 李新江、叶保华两位老师协助采集标本表示 谢意。

REFERENCES

- Bei-Bienko, G. J. (Ya.) and Mishchenko, L. L. 1951. Acridoidea of the fauna of the USSR and neighbouring countries. Pts. 2 Opred. Faune SSSR, Moscow. pp. 543-545.
- Bi, D-Y and Xia, K-L 1987. Agricultural Insects, Plant Diseases and Weed of Xizang. Vol. 1. 58 - 59, figs 15 - 21.
- Chang, F-S 1937. Some new Acridids from Szechwan and Szechwan-Tibetan border (Orthoptera: Acrididae). Notes Ent. Chinoise, Mus. Heude, Shanghai, 4 (8): 193-194, pl. IV. fig. 9.
- Cheng, T-M and Chiu, C-H 1965. A survey of grasshoppers from Ningsia and Northern Shensi. *Acta Entomologica Sinica*, 14 (6): 585 587, figs 2 8.
- Eades, D. C., Otte, D., Cigliano, M. M. and Braun, H. Orthoptera Species File Online. Version 2. 0/4. 0. [2011. 2]. < http:// Orthoptera. Species File. org >.
- Fischer-Waldheim, 1846. Orhtoptera Ress. N. 23. 305, tab. 23, fig. 4.
- Harz, K. 1975. Orthoptera Europe, II. 916 918.
- Ikonnikov, N. 1913. Uber die von P. Schmidt aus Korea Mifgebrachten Acridoideen; 15.
- Jin, X-B and Zhang, F-L 1983. A new species of the genus Euchorthippus Tarbinsky from China (Orthoptera: Acridoidea). Zoological Research, 4 (4): 377 – 381, figs 1 – 12.
- Liang, G-Q and Jia, F-L 1992. One new genus and four new species of grasshoppers from Inner Mongolia, China (Orthoptera: Acridoidea). Acta Scientiarum Naturalium Universitatis Sunyatseni, 31 (1): 93-98, figs 17-21.
- Liu, J-P 1990. A handbook for identifying grasshoppers of China. Tianze Press Xian, pp. 155 – 157.
- Ren, B-Z and Zhao, Z 2001. A new species of genus Euchorthippus Tarbinsky from Jilin Province (Orthoptera: Acridoidea). Acta Zootaxonomica Sinica, 26 (2): 191-193, figs 1-6. [动物分类学规]
- Tarbinsky, S. P. 1925. Russkoe entomologicheskoe obozrenie. 19: 192.
- Tu, Chin and Cheng, T-M 1964. A study of the family Acrididae of

- Shensi II. Genus *Euchorthippus* Tarbinsky, 1925 and genus *Mongolotettix* Rehn, 1928. *Acta Zoologica Sinica*, 16 (3): 474 476, figs 1 3.
- Xia, K-L 1958. Syn Classfication Chinese Acridoidea. 129.
- Yin, X-C 1984. Grasshoppers and locusts from Qinghai-Xizang Plateau. Science Press, Beijing. pp. 159 – 160, figs 311 – 313. [In Chinese with English summary]
- Yin, X-C, Shi, J-P and Yin, Z 1996. A Synonymic Catalogue of Grasshoppers and Their Allies of the World. China Forestry Publishing House, Beijing. pp. 270 – 272.
- Zhang, F-L and Jin, X-B 1985. A preliminary survey of grasshoppers from Da Xinganling, Northeast China (Orthoptera: Acridoidea). Contr. Shanghai Institue Entomology, 5: 216 – 217, figs 38 – 44.
- Zhang, F-L and Ren, B-Z 1992. A new species of gemus *Euchorthippus* Tarbinsky from China (Orthoptera: Acridoidea). *Entomotaxonomia*, 14 (1): 11-13, figs 1-5.
- Zhang, F-L and Ren, B-Z 1992. A new species of genus Euchorthippus Tarbinsky from Jiling Province (Orthoptera: Acridoidea). Acta Entomologica Sinica, 35 (4): 456-458, figs 1-5.
- Zheng, Z-M 1980a. A new species of grasshoppers from Shaanxi Province. *Entomotaxonomia*, 2 (1): 75 76, figs 1 3.
- Zheng, Z-M 1980b. New genera and species of grasshoppers from Sichuan, Shaanxi and Yuuman. *Entomotaxonomia*, 2 (4): 344 345, figs 49 57.
- Zheng, Z-M 1985. Acridoidea from YGSSN Regions. 327.
- Zheng, Z-M 1993. Taxonomy of Acrididae. Shaanxi Normal University Press, Xi'an, China. pp. 335 – 343; figs 1 075 – 1 099.
- Zheng, Z-M and He, D-H 1993. Two new species of Arcypteridae from Ningxia. Journal Ningxia Agricultural College, 14 (4): 7 9, figs 1-5.
- Zheng, Z-M and Lu, R-S 2002. Two new species of grasshoppers from Zhontiaoshan, Shanxi Province (Orthoptera: Acridoidea). Entomotaxonomia, 24 (1): 16-20, figs 4-7.
- Zheng, Z-M and Xia, K-L 1998. Fauna Sinica Insecta, Vol. 10, Orthoptera, Acridoidea, Oedipodidae and Arcypteridae, Beijing. pp. 489 – 505, figs 290 – 301.
- Zheng, Z-M and Wang, X-R 1993. Two new species of genus Euchorthippus Tarbinsky from Shanxi Province (Orthoptera: Arcypteridae). Sichuan Journal of Zoology, 12 (2): 4-7, figs 1-8.